

On Low-Energy Effective Action of Noncommutative Hypermultiplet Model

I.B. Samsonov¹

Department of Quantum Field Theory, Tomsk State University, Tomsk 634050, Russia

Abstract

We consider the noncommutative hypermultiplet model within harmonic superspace approach. The 1-loop four-point contributions to the effective action of selfinteracting q -hypermultiplet are computed. This model has two coupling constants instead of a single one in commutative case. It is shown that both these coupling constants are generated by 1-loop quantum corrections in the model of q -hypermultiplet interacting with vector multiplet. The holomorphic effective action of q -hypermultiplet in external gauge superfield is calculated. For the fundamental representation there is no UV/IR-mixing and the holomorphic potential is a $\star$ -product generalization of a standard commutative one. For the adjoint representation of $U(N)$ gauge group the leading contributions to the holomorphic effective action are given by the terms respecting for the UV/IR-mixing which are related to $U(1)$ phase of $U(N)$ group.

Noncommutative field theory attracts much attention due to its remarkable properties and profound links with modern string/brane activity. This theory is based on an idea of underlying noncommutative space-time geometry [1, 2] that allows to incorporate a nonlocal structure of the theory and a possibility to apply the standard quantum field theory methods (see ref. [3] for review).

Noncommutative field theories possess specific properties of UV- and IR-divergences (the UV/IR-mixing [4]) which distinguish them from ordinary theories very essentially. For example, the insertion of noncommutativity can break the renormalizability of field models [5]. The presence of nonplanar diagrams in gauge theories [11, 15, 16] leads to the deformation of $\star$ -product by quantum corrections ($\star_n$ -products appear) that breaks the manifest gauge symmetry of effective action. The restoration of manifest gauge invariance of effective action involves the open Wilson line operators [15, 16] which absorb all $\star_n$ -products into gauge invariant objects.

Most of these problems are convenient to explore in the frame of supersymmetric theories. The gauge invariance of effective action had been examined by many authors on the example of noncommutative $\mathcal{N} = 4$ SYM model [11, 15] since this model is finite and have no UV/IR-mixing [14]. The structure of low-energy effective action in the model of $\mathcal{N} = 2$ SYM was

¹E-mail: samsonov@phys.tsu.ru

studied in the papers [8, 9, 10, 11, 12], effective potential in noncommutative Wess-Zumino model was considered in [13].

Our aim is to investigate the low-energy effective action of $\mathcal{N} = 2$ supersymmetric model of matter fields (hypermultiplet model) on noncommutative plain. A natural approach for studying the $\mathcal{N} = 2$ supersymmetric models is a harmonic superspace method [18]. It provides an explicit $\mathcal{N} = 2$ supersymmetry at all steps of quantum computations. The $\mathcal{N} = 2$ harmonic superspace is obtained from the conventional $\mathcal{N} = 2$ superspace by adding the harmonic variables which parameterize the $SU(2)/U(1)$ coset of internal automorphisms group of $\mathcal{N} = 2$ supersymmetry algebra (see [19] for details). The generalization of this construction to the case of noncommutative geometry is straightforward. We suppose that only the bosonic coordinates of superspace obey the Moyal-Weyl commutation relations

$$[x_\mu, x_\nu] = i\theta_{\mu\nu}, \quad (1)$$

where $\theta_{\mu\nu}$ is the constant antisymmetric tensor while the fermionic coordinates of superspace are still anticommuting. Such a deformation does not break supersymmetry since the derivatives ∂_μ are supercovariant. This is just the simplest way to introduce the noncommutativity without breaking of supersymmetry. Note however some attempts to construct a superspace with non-anticommuting Grassman coordinates [7] where the algebraic aspects of such deformations were studied. A general case of deformations of superspace was considered also in [6]. The harmonic variables remain commutative because they correspond to the $SU(2)$ automorphisms algebra which is not deformed since the deformations in the bosonic sector only do not modify the supersymmetry algebra [6, 7]. The noncommutative $\mathcal{N} = 2$ harmonic superspace was considered firstly in [17].

A natural multiplication of superfields on noncommutative plain (1) is given by the $\star$ -product

$$(\phi \star \psi)(x) = \phi(x) e^{\frac{i}{2}\theta^{\mu\nu} \overleftarrow{\partial}_\mu \overrightarrow{\partial}_\nu} \psi(x), \quad (2)$$

which is associative but noncommutative.

In this letter we consider the noncommutative models of selfinteracting q -hypermultiplet and q -hypermultiplet interacting with vector superfield. In commutative case these theories were studied in [18]-[26]. We are interested mainly in new features of these theories appearing due to the noncommutativity. The model of selfinteracting commutative q -hypermultiplet is known to be nonrenormalizable [18], but it was shown that the selfinteraction can be induced by quantum corrections from the model of q -hypermultiplet interacting with vector multiplet [23]. For the model of q -hypermultiplet in external gauge superfield the low-energy effective action is defined by the holomorphic potential which was computed exactly in papers [24, 25] using the harmonic superspace technique. In this paper we show how these results can be generalized to noncommutative case.

To illustrate the features of noncommutative quantum field theory in harmonic superspace we consider the simplest q -hypermultiplet model with the quartic selfinteraction of the type $\lambda(\tilde{q}^+)^2(q^+)^2$ [18]. The action of corresponding noncommutative model is obtained by insertion of $\star$ -product (2) instead of usual multiplication ¹

$$S[q, \tilde{q}] = S_0[q, \tilde{q}] + S_{int}[q, \tilde{q}], \quad (3)$$

¹ We follow the notations for harmonic superspace objects which were accepted throughout the papers [22]-[26].

$$S_0[q, \check{q}] = \int d\zeta^{(-4)} \check{q}^+ D^{++} q^+, \quad (4)$$

$$S_{int}[q, \check{q}] = \int d\zeta^{(-4)} (\alpha \check{q}^+ \star \check{q}^+ \star q^+ \star q^+ + \beta \check{q}^+ \star q^+ \star \check{q}^+ \star q^+). \quad (5)$$

Note that there is a possibility to introduce the two coupling constants α, β instead of a single one λ in commutative case due to two types of ordering of superfields. This situation is similar to noncommutative ϕ^4 model [5]. In commutative limit $\theta \rightarrow 0$ the two terms in the interaction (5) reduce to a single one $\lambda(q^+)^2(\check{q}^+)^2$.

The propagator corresponding to the free action (4) is given by [18]

$$\langle \check{q}(\zeta_1) q(\zeta_2) \rangle = G_0^{(1,1)}(1|2) = -\frac{1}{\square_1} \frac{(D_1^+)^4 (D_2^+)^4 \delta^{12}(z_1 - z_2)}{(u_1^+ u_2^+)^3}. \quad (6)$$

The action S_{int} (5) defines the four-point vertex (in momentum space)

$$(2\pi)^4 \delta^4(p_1 + p_2 + p_3 + p_4) [\alpha \cos \frac{p_1 \theta p_2}{2} \cos \frac{p_3 \theta p_4}{2} + \beta \cos(\frac{p_1 \theta p_3}{2} + \frac{p_2 \theta p_4}{2})]. \quad (7)$$

We consider also the model of massive q -hypermultiplet with the free action

$$S_0[q, \check{q}] = \int d\zeta^{(-4)} \check{q}^+ (D^{++} + iV_0^{++}) q^+ \equiv \int d\zeta^{(-4)} \check{q}^+ \mathcal{D}^{++} q^+, \quad (8)$$

where $V_0^{++} = -\bar{W}_0(\theta^+)^2 - W_0(\bar{\theta}^+)^2$, $\bar{W}_0 W_0 = m^2$ is the mass. The corresponding propagator was found in refs. [20, 21, 23] in the form

$$\langle \check{q}(\zeta_1) q(\zeta_2) \rangle = G^{(1,1)}(1|2) = -\frac{1}{\square_1 + m^2} (D_1^+)^4 (D_2^+)^4 \left\{ \frac{e^{i\Omega_0(1) - i\Omega_0(2)} \delta^{12}(z_1 - z_2)}{(u_1^+ u_2^+)^3} \right\}, \quad (9)$$

where $\Omega_0 = -\bar{W}_0 \theta^+ \theta^- - W_0 \bar{\theta}^+ \bar{\theta}^-$ is a bridge superfield [23].

An interaction of q -hypermultiplet (8) with the vector multiplet is introduced by minimal way $\mathcal{D}^{++} \rightarrow \mathcal{D}^{++} + iV^{++} \star$. For the fundamental representation of $U(1)$ gauge group it reads

$$S_{int,f} = \int d\zeta^{(-4)} \check{q}^+ \star iV^{++} \star q^+. \quad (10)$$

The corresponding vertex is given by

$$(2\pi)^4 \delta^4(p + k_1 + k_2) e^{-\frac{i}{2} k_1 \theta k_2}. \quad (11)$$

The q -hypermultiplet model (8) with the interaction (10) is invariant under the following gauge transformations

$$\begin{aligned} \delta q^+ &= i\lambda \star q^+, & \delta \check{q}^+ &= -i\check{q}^+ \star \lambda, \\ \delta V^{++} &= -D^{++} \lambda + i\lambda \star V^{++} - iV^{++} \star \lambda. \end{aligned} \quad (12)$$

A generalization to the case of non-Abelian gauge group is trivial: the vector superfield V^{++} should be a matrix of $U(N)$ group while the hypermultiplets q^+ , $\check{q}^+$ belong to the fundamental representation.

It is interesting to study also the adjoint representation of gauge group when the interaction is given by

$$S_{int,ad} = i \text{tr} \int d\zeta^{(-4)} \check{Q}^+ \star [V^{++}, Q^+]_{\star}, \quad (13)$$

where $[V^{++}, Q^+]_{\star} = V^{++} \star Q^+ - Q^+ \star V^{++}$ and the superfields V^{++} , Q^+ , $\check{Q}^+$ are the matrices of $U(N)$ group. For the case of adjoint representation the gauge transformations look like

$$\begin{aligned} \delta Q^+ &= i[\lambda, Q^+]_{\star}, & \delta \check{Q}^+ &= -i[\check{Q}^+, \lambda]_{\star}, \\ \delta V^{++} &= -D^{++} \lambda + i[\lambda, V^{++}]_{\star}. \end{aligned} \quad (14)$$

Now we begin with the quantum computations of 1-loop four-point diagram in the model (3)

$$\begin{aligned} \Gamma_4^{(1)}[\check{q}^+, q^+] &= \Gamma_s[\check{q}^+, q^+] + \Gamma_t[\check{q}^+, q^+], \\ \Gamma_s[\check{q}^+, q^+] &= \text{diagram with two vertices, external lines } q^+(p_1), q^+(p_2), q^+(p_3), q^+(p_4), \text{ and internal lines } k, p_1+p_3-k, \end{aligned} \quad (15)$$

Using the Feynman rules (6,7) we obtain the following expressions for these diagrams

$$\begin{aligned} \Gamma_{s,t} &= \int \frac{d^4 p_1 \dots d^4 p_4}{(2\pi)^{16}} d^4 \theta_1^+ d^4 \theta_2^+ du_1 du_2 \delta^4(p_1 + p_2 + p_3 + p_4) \check{q}^+(1) \check{q}^+(2) q^+(3) q^+(4) \\ &\times \frac{(D_1^+)^4 (D_2^+)^4}{(u_1^+ u_2^+)^3} \delta^8(\theta_1 - \theta_2) \frac{(D_1^+)^4 (D_2^+)^4}{(u_1^+ u_2^+)^3} \delta^8(\theta_1 - \theta_2) I_{s,t}(p_1, \dots, p_4), \end{aligned} \quad (16)$$

where

$$\begin{aligned} I_t(p_1, \dots, p_4) &= 2 \int d^4 k \frac{F_t(p_1, \dots, p_4, k)}{k^2 (p_1 + p_2 - k)^2}, \\ I_s(p_1, \dots, p_4) &= 8 \int d^4 k \frac{F_s(p_1, \dots, p_4, k)}{k^2 (p_1 + p_3 - k)^2}, \end{aligned} \quad (17)$$

$F_{s,t}(p_1, \dots, p_4, k)$ are the functions the structure of which is stipulated by noncommutativity

$$\begin{aligned} F_t(p_1, \dots, p_4, k) &= (\alpha \cos \frac{p_1 \theta p_2}{2} \cos \frac{k \theta (p_1 + p_2)}{2} + \beta \cos(\frac{p_1 \theta k}{2} + \frac{p_2 \theta (p_1 - k)}{2})) \\ &\times (\alpha \cos \frac{k \theta (p_1 + p_2)}{2} \cos \frac{p_3 \theta p_4}{2} + \beta \cos(\frac{k \theta p_3}{2} - \frac{(p_3 + k) \theta p_4}{2})), \\ F_s(p_1, \dots, p_4, k) &= (\alpha \cos \frac{k \theta p_1}{2} \cos \frac{(p_1 - k) \theta p_3}{2} + \beta \cos(\frac{k \theta (p_1 + p_3)}{2} + \frac{p_1 \theta p_3}{2})) \\ &\times (\alpha \cos \frac{k \theta p_4}{2} \cos \frac{(p_4 + k) \theta p_2}{2} + \beta \cos(\frac{p_2 \theta p_4}{2} - \frac{k \theta (p_1 + p_3)}{2})). \end{aligned} \quad (18)$$

Further computations are very similar to usual ones in commutative q^4 -model given in [18]. The resulting expression for the effective action (16) reads

$$\Gamma_{s,t} = \int \frac{d^4 p_1 \dots d^4 p_4}{(2\pi)^{16}} d^8 \theta du_1 du_2 \frac{\check{q}^+(1) \check{q}^+(2) q^+(3) q^+(4)}{(u_1^+ u_2^+)^2} \delta^4(p_1 + p_2 + p_3 + p_4) I_{s,t}(p_1, \dots, p_4). \quad (19)$$

This effective action differs from the corresponding commutative one by the presence of two functions $F_{s,t}$ in the integrals (17). These functions change the UV-structure of integrals $I_{s,t}$ and lead to the appearance of UV/IR-mixing.

To study the structure of momentum integrals (17) let us split the functions $F_{s,t}$ into planar (pl) and non-planar (npl) parts

$$F_{s,t}(p_1, \dots, p_4, k) = F_{s,t}^{pl}(p_1, \dots, p_4) + F_{s,t}^{npl}(p_1, \dots, p_4, k), \quad (20)$$

where we assume that $F_{s,t}^{pl}$ do not contain the terms depending on the momentum k . The planar parts are written explicitly

$$\begin{aligned} F_t^{pl} &= \frac{\alpha^2}{2} \cos \frac{p_1 \theta p_2}{2} \cos \frac{p_3 \theta p_4}{2}, \\ F_s^{pl} &= \left(\frac{\alpha^2}{8} + \frac{\beta^2}{2} \right) \cos \left(\frac{p_1 \theta p_3}{2} + \frac{p_2 \theta p_4}{2} \right) + \frac{\alpha \beta}{2} \cos \left(-\frac{p_1 \theta p_3}{2} + \frac{p_2 \theta p_4}{2} \right). \end{aligned} \quad (21)$$

Substituting the expressions (21) into integrals (17)

$$\begin{aligned} I_t^{pl}(p_1, \dots, p_4) &= 2F_t^{pl}(p_1, \dots, p_4) \int \frac{d^4 k}{k^2(p_1 + p_2 - k)^2}, \\ I_s^{pl}(p_1, \dots, p_4) &= 8F_s^{pl}(p_1, \dots, p_4) \int \frac{d^4 k}{k^2(p_1 + p_3 - k)^2}, \end{aligned} \quad (22)$$

one can see that the planar contributions have the same divergences as corresponding diagrams of commutative q -hypermultiplet. The momentum integrals in eq. (22) have the IR-divergences at low external momenta which do not appear in massive theory (these singularities are not related to the UV/IR-mixing) and UV-divergences which can not be renormalized since the coupling constants α, β have the mass dimension -2 as in commutative theory [18]. So, the selfinteracting noncommutative q -hypermultiplet model is nonrenormalizable as well as the corresponding commutative theory. Inserting the noncommutativity does not improve the situation with renormalizability here.

Let us study now the structure of non-planar diagrams where the effect of UV/IR-mixing arises. One can show that all non-planar terms of functions (18) look like

$$\cos \frac{p_1 \theta p_2}{2} \cos \frac{p_3 \theta p_4}{2} \cos(k\theta(p_1 + p_2))$$

with various combinations of external momenta. They define the structure of momentum integrals of non-planar type

$$\begin{aligned} I_t^{npl}(p_1, \dots, p_4) &\sim \cos \frac{p_1 \theta p_2}{2} \cos \frac{p_3 \theta p_4}{2} \int d^4 k \frac{e^{ik\theta(p_1+p_2)}}{k^2(p_1 + p_2 - k)^2} \\ &= \cos \frac{p_1 \theta p_2}{2} \cos \frac{p_3 \theta p_4}{2} \int_0^1 d\xi K_0(\sqrt{\xi(1-\xi)P}), \end{aligned} \quad (23)$$

where $P = (p_1 + p_2)^2 \cdot (p_1 + p_2) \circ (p_1 + p_2)$, $p_1 \circ p_2 = p_1^\mu(\theta\theta)_{\mu\nu}p_2^\nu$, K_0 is the modified Bessel function. This expression has no UV-divergence but it is singular at low momenta $p_i \rightarrow 0$ due to the asymptotics of Bessel function

$$K_0(x) \xrightarrow{x \rightarrow 0} -\ln \frac{x}{2} + \text{finite}. \quad (24)$$

Therefore at low external momenta the integral over momentum k behaves as

$$\int d^4 k \frac{e^{ik\theta(p_1+p_2)}}{k^2(p_1 + p_2 - k)^2} \sim \ln \frac{1}{P}. \quad (25)$$

The expression (25) is singular in commutative limit $\theta \rightarrow 0$. Such a singularity of effective action was called the UV/IR-mixing [4]. Therefore the effective action of noncommutative hypermultiplet does not reduce to a standard one in commutative limit.

As a result, the model of noncommutative hypermultiplet is nonrenormalizable and has the UV/IR-mixing in the sector of non-planar diagrams.

To obtain a renormalizable model in commutative case it was suggested [23] to consider the model (8) and generate the interaction like (5) by 1-loop low-energy corrections to the effective action of the type

$$\Gamma_4 = \Gamma_s + \Gamma_t, \quad \Gamma_s = \text{diagram}, \quad \Gamma_t = \text{diagram} \quad (26)$$

We study the analogical possibility in noncommutative hypermultiplet model. In the low energy approximation the loops shrink down to points and we will see that the finite parts of these diagrams give the coupling constants α, β of selfinteracting q -hypermultiplet (5) induced only by quantum corrections. Induced selfinteraction in the commutative hypermultiplet model was considered in [23].

To compute these diagrams we apply the Feynman rules obtained (9,11) and the following expression for the propagator of intermediate gauge boson [18]

$$i < V^{++}(1)V^{++}(2) > = \frac{1}{\square_1} (D_1^+)^4 \delta^{12}(z_1 - z_2) \delta^{(-2,2)}(u_1, u_2). \quad (27)$$

Using the explicit expressions for the propagators and vertices one obtains the following representations for the diagrams (26)

$$\Gamma_{s,t}[\check{q}^+, q^+] = -\frac{iq^4}{(2\pi)^{16}} \int d^4 p_1 d^4 p_2 d^4 p_3 d^4 p_4 F_{s,t}(p_1, \dots, p_4) I_{s,t}(p_1, \dots, p_4) \times \int d^8 \theta \frac{du_1 du_2}{(u_1^+ u_2^+)^2} q^+(p_1) q^+(p_2) \check{q}^+(p_3) \check{q}^+(p_4) e^{i\Omega_0(2) - i\Omega_0(1)} e^{i\Omega_0(2) - i\Omega_0(1)}, \quad (28)$$

where

$$I_s(p_1, \dots, p_4) = \frac{1}{(2\pi)^4} \int \frac{d^4 k}{k^2 (k + p_1 + p_3)^2} \frac{1}{((k + p_1)^2 + m^2)((k - p_4)^2 + m^2)}, \quad (29)$$

$$I_t(p_1, \dots, p_4) = \frac{1}{(2\pi)^2} \int \frac{d^4 k e^{ik\theta(p_1 + p_3)}}{(k + p_3)^2 (k - p_1)^2} \frac{1}{(k^2 + m^2)((k + p_3 + p_4)^2 + m^2)},$$

$$F_s(p_1, \dots, p_4) = e^{-\frac{i}{2} p_1 \theta p_3} e^{-\frac{i}{2} p_2 \theta p_4}, \quad F_t(p_1, \dots, p_4) = e^{-\frac{i}{2} p_2 \theta p_1} e^{-\frac{i}{2} p_3 \theta p_4}. \quad (30)$$

It should be noted that the momentum integrals (29) are UV-finite. Therefore there is no UV/IR-mixing and these diagrams are well defined in the limits $\theta \rightarrow 0$ and $p_i \rightarrow 0$. At zero external momenta both these integrals are computed exactly:

$$I_{s,t} = \frac{1}{(4\pi)^2 m^2} \left(\frac{1}{m^2} \ln\left(1 + \frac{m^2}{\Lambda_{s,t}^2}\right) - \frac{1}{\Lambda_{s,t}^2 + m^2} \right), \quad (31)$$

where $\Lambda_{s,t}$ are the parameters of infrared cutoff which do not coincide generally speaking. The values of $\Lambda_{s,t}$ should be taken less than the mass of the lightest particle that corresponds to the Wilsonian low-energy effective action.

In order to extract the relevant low-energy contribution from eq. (28) it is necessary to employ the covariant derivatives algebra [18]

$$[\mathcal{D}^{++}, \mathcal{D}^{--}] = D^0, \quad (32)$$

where

$$\mathcal{D}^{\pm\pm} = D^{\pm\pm} + iV_0^{\pm\pm}, \quad V_0^{\pm\pm} = D^{\pm\pm}\Omega_0. \quad (33)$$

The relation (32) is used to prove the identity

$$\begin{aligned} q^+ \star q^+ \star \check{q}^+ \star \check{q}^+ e^{i\Omega_0(2)-i\Omega_0(1)} e^{i\Omega_0(2)-i\Omega_0(1)} \\ = \frac{1}{2}[\mathcal{D}_1^{++}, \mathcal{D}_1^{--}] q^+ \star q^+ \star \check{q}^+ \star \check{q}^+ e^{i\Omega_0(2)-i\Omega_0(1)} e^{i\Omega_0(2)-i\Omega_0(1)}. \end{aligned} \quad (34)$$

Then it is easy to argue that only the first term $\sim \mathcal{D}^{++}\mathcal{D}^{--}$ in this identity gives the leading low-energy contribution. Integrating by parts one can cancel the harmonic distribution $\frac{1}{(u_1^+ u_2^+)^2}$ by the use of another identity [18]

$$D_1^{++} \frac{1}{(u_1^+ u_2^+)^2} = D_1^{--} \delta^{(2,-2)}(u_1, u_2). \quad (35)$$

The harmonic δ -function on the right hand side of eq. (35) removes one of the two remaining harmonic integrals and allows us to rewrite eq. (28) to the form

$$\Gamma_{s,t}[\check{q}^+, q^+] = -ig^4 I_{s,t} \cdot S_{s,t}[\check{q}^+, q^+], \quad (36)$$

where

$$\begin{aligned} S_s &= \int d^4x d^8\theta du \mathcal{D}^{--} \mathcal{D}^{--} q^+ \star \check{q}^+ \star q^+ \star \check{q}^+, \\ S_t &= \int d^4x d^8\theta du \mathcal{D}^{--} \mathcal{D}^{--} q^+ \star q^+ \star \check{q}^+ \star \check{q}^+. \end{aligned} \quad (37)$$

The integrals over full $\mathcal{N} = 2$ superspace in eqs. (37) can be transformed to ones over the analytic subspace

$$\begin{aligned} S_s &= -2m^2 \int d\zeta^{(-4)} q^+ \star \check{q}^+ \star q^+ \star \check{q}^+, \\ S_t &= -2m^2 \int d\zeta^{(-4)} q^+ \star q^+ \star \check{q}^+ \star \check{q}^+, \end{aligned} \quad (38)$$

where we have used the explicit form for v^{--} given by eq. (33) and relation $\bar{W}W = m^2$. As a result, the low-energy part of the effective action (36) is represented in the form of four-point q -hypermultiplet interaction (8)

$$\Gamma_4 = \int d\zeta^{(-4)} (\lambda_t \check{q}^+ \star \check{q}^+ \star q^+ \star q^+ + \lambda_s \check{q}^+ \star q^+ \star \check{q}^+ \star q^+), \quad (39)$$

where the coupling constants $\lambda_{s,t}$ appear due to quantum corrections (31) as

$$\lambda_{s,t} = \frac{g^2}{(4\pi)^2} \left(\frac{1}{m^2} \ln(1 + \frac{m^2}{\Lambda_{s,t}}) - \frac{1}{\Lambda_{s,t}^2 + m^2} \right). \quad (40)$$

Note that the effective action (39) is smooth in the limit $\theta \rightarrow 0$ and both induced coupling constants $\lambda_{s,t}$ (40) reduce to a single coupling constant obtained in ref. [23] for the commutative hypermultiplet model.

Let us consider now the low-energy effective action of massive single q -hypermultiplet model (8) in fundamental representation of $U(1)$ gauge group. The 1-loop effective action in Coulomb branch is given by

$$\Gamma^{(1)}[V^{++}] = i \text{tr} \ln(\nabla^{++} \star) = \sum_{n=1}^{\infty} \Gamma_n, \quad (41)$$

$$\Gamma_n = i \frac{(-i)^n}{n} \int d\zeta_1^{(-4)} \dots d\zeta_n^{(-4)} G^{(1,1)}(\zeta_1, \zeta_2) \star V^{++}(\zeta_2) \dots G^{(1,1)}(\zeta_n, \zeta_1) \star V^{++}(\zeta_1),$$

where $\nabla^{++} \star = D^{++} + iV_0^{++} + iV^{++} \star$. In momentum space the n -point function Γ_n reads

$$\Gamma_n = \frac{i(-i)^n}{n(2\pi)^{4n}} \int d^4 p_1 d^4 \theta_1^+ du_1 \dots d^4 p_n d^4 \theta_n^+ du_n \int d^4 k \delta^4(\sum p_i) e^{-\frac{i}{2} k \theta (\sum p_i)} \quad (42)$$

$$\times G^{(1,1)}(k + p_2 + p_3 + \dots + p_n) G^{(1,1)}(k + p_3 + \dots + p_n) \dots G^{(1,1)}(k + p_n) G^{(1,1)}(k) \\ \times e^{-\frac{i}{2} \sum_{i < j}^n p_i \theta p_j} V^{++}(1) \dots V^{++}(n).$$

The first exponent in eq. (42) reduces to unity due to the relation

$$e^{-\frac{i}{2} k \theta (\sum p_i)} \delta^4(\sum p_i) = \delta^4(\sum p_i). \quad (43)$$

The equation (43) ensures the absence of nonplanar diagrams and UV/IR-mixing. Therefore, the limit $\theta \rightarrow 0$ is smooth and in low-energy approximation (when we neglect all derivatives of V^{++}) the factor $e^{-\frac{i}{2} \sum_{i < j}^n p_i \theta p_j}$ can be dropped. In such an approximation the effective action of noncommutative q -hypermultiplet reduces to a standard one of commutative q -hypermultiplet calculated in ref. [25]. A non-trivial result here is the absence of non-planar diagrams and UV/IR-mixing.

Now consider the next order in the approximation when all derivatives related to non-commutativity are kept. In this case we keep the factor $e^{-\frac{i}{2} \sum_{i < j}^n p_i \theta p_j}$ in eq. (42) and use it to restore the $\star$ -product of superfields due to the identity

$$\int d^4 p_1 \dots d^4 p_n e^{-\frac{i}{2} \sum_{i < j}^n p_i \theta p_j} V_1(p_1) \dots V_n^{++}(p_n) \delta^4(\sum p_i) \quad (44)$$

$$= \int d^4 x V_1^{++}(x) \star V_2^{++}(x) \star \dots \star V_n^{++}(x).$$

We expect the resulting expression for the effective action to be gauge invariant. It means that we have to express the effective action in terms of strength superfield which is a nonlinear function of V^{++} even for the Abelian gauge group (it can be found in complete analogy with commutative case [21])

$$W = \frac{1}{4} \bar{D}_{\dot{\alpha}}^+ \bar{D}^{+\dot{\alpha}} \sum_{n=1}^{\infty} i(-ig)^n \int du_1 \dots du_n \frac{V^{++}(z, u_1) \star V^{++}(z, u_2) \star \dots \star V^{++}(z, u_n)}{(u^+ u_1^+)(u_1^+ u_2^+) \dots (u_n^+ u^+)}. \quad (45)$$

To simplify this problem we choose the external gauge superfield in the form

$$\tilde{W} = -\frac{1}{4} \int du \bar{D}_{\dot{\alpha}}^- \bar{D}^{-\dot{\alpha}} \tilde{V}^{++}(z, u), \quad (46)$$

that corresponds to the first term in series (45). With such a special background $\tilde{W}$ one obtains the standard expression for holomorphic effective action [25] written in terms of

strength superfields $\tilde{W}$ where conventional multiplication should be replaced by $\star$ -product due to the relation (44). To return to arbitrary background it is necessary to restore the full strength W from $\tilde{W}$ with the help of gauge transformations

$$W = e_{\star}^{i\lambda} \star \tilde{W} \star e_{\star}^{-i\lambda}$$

with a special $\lambda = \lambda(W)$. Therefore, for arbitrary W the holomorphic effective action of noncommutative q -hypermultiplet reads

$$\Gamma[V^{++}] = -\frac{1}{64\pi^2} \int d^4x d^4\theta W \star W \star \ln_{\star} \frac{W}{\Lambda} + c.c. \quad (47)$$

Note that this expression is manifestly gauge invariant and has correct commutative limit.

A generalization of this result to the case of $U(N)$ gauge group broken down to $[U(1)]^N$ is rather trivial since the strength W belonging to $[U(1)]^N$ can be chosen as $W = \text{diag}(W_1, \dots, W_N)$. Therefore the effective action of such theory is a sum of actions (47)

$$\Gamma_{[U(1)]^N}[V^{++}] = \sum_{i=1}^N \Gamma_{U(1)}[V_i^{++}], \quad (48)$$

where $\Gamma_{U(1)}[V_i^{++}]$ is given by eq. (47).

In the case of adjoint representation of gauge group the situation is more complicated. Let us consider the classical action of q -hypermultiplet in adjoint representation when the vector superfield belongs to the Cartan subalgebra of $u(N)$, i.e.

$$Q^+ = \sum_{i,j}^N q_{ij}^+ e_{ij}, \quad \check{Q}^+ = \sum_{i,j}^N \check{q}_{ij}^+ e_{ji}, \quad V^{++} = \sum_{k=1}^N V_k^{++} e_{kk}, \quad (49)$$

where

$$(e_{ij})_{kl} = \delta_{ik} \delta_{jl}, \quad i, j, k, l = 1, \dots, n \quad (50)$$

is a Cartan-Weyl basis of $u(N)$ algebra [29]. One can easily check that the interaction (13) now looks like

$$\text{tr} \int d\zeta^{(-4)} \check{Q}^+ \star [V^{++}, Q^+]_{\star} = \sum_{k,l}^N \int d\zeta^{(-4)} (\check{q}_{kl}^+ \star V_k^{++} \star q_{kl}^+ - \check{q}_{kl}^+ \star q_{kl}^+ \star V_l^{++}). \quad (51)$$

Therefore the contributions from hypermultiplets q_{ij}^+ , q_k^+ factorize and the formal expression for the 1-loop effective action reads

$$\Gamma[V^{++}] = \sum_{k,l=1}^N \text{Tr} \ln \nabla_{kl}^{++}. \quad (52)$$

where the operators ∇_{kl}^{++} , act on hypermultiplets by the rule

$$\nabla_{kl}^{++} q_{kl}^+ = \mathcal{D}^{++} q_{kl}^+ + i V_k^{++} \star q_{kl}^+ - i q_{kl}^+ \star V_l^{++}.$$

The effective action (52), unlike the fundamental representation, contain the terms respecting for the UV/IR-mixing which appear in two-point diagrams. These diagrams can be computed

in complete analogy as it was done in the case of fundamental representation. The resulting expression for the nonplanar part of two-point one-loop effective action reads

$$\Gamma_2^{np}[V^{++}] = \frac{1}{16\pi^2} \int d^4p d^8\theta du_1 du_2 \int_0^1 d\xi K_0(\sqrt{(\xi(1-\xi)p^2 + m^2)p \circ p}) \times \frac{\mathcal{V}^{++}(p, \theta, u_1) \mathcal{V}^{++}(-p, \theta, u_2)}{(u_1^+ u_2^+)^2}, \quad (53)$$

where $\mathcal{V}^{++} = \sum_{i=1}^N V_i^{++}$ is the superfield corresponding to $U(1)$ subgroup of $U(N)$ gauge group. In the low-energy approximation we employ the asymptotics of the Bessel function (24) to extract the part of the effective action which is singular in the commutative limit $\theta \rightarrow 0$

$$\Gamma_2^{np}[V^{++}] = \frac{1}{32\pi^2} \int d^4p d^8\theta du_1 du_2 \ln \frac{4}{m^2 p \circ p} \frac{\mathcal{V}^{++}(p, \theta, u_1) \mathcal{V}^{++}(-p, \theta, u_2)}{(u_1^+ u_2^+)^2}. \quad (54)$$

We see that the UV/IR-mixing appears only in the $U(1)$ sector that was firstly observed for noncommutative $U(N)$ Yang-Mills model in [8] and for supersymmetric gauge theories in [9, 10].

The next order contribution to the effective action which is finite in the limit $\theta \rightarrow 0$ corresponds to the holomorphic potential of commutative q -hypermultiplet in adjoint representation of $SU(N)$ group calculated in [26].

Let us summarize the results. We have considered the model of noncommutative selfinteracting hypermultiplet using the harmonic superspace approach. It is worth pointing out that such a model has two coupling constants instead of a single one in commutative case. The 1-loop four-point contributions to the effective action of selfinteracting q -hypermultiplet are calculated. The ultraviolet divergences in this model are responsible for the appearance of UV/IR-mixing terms. These divergences can not be renormalized as well as in commutative case. We find that both these coupling constants are generated by 1-loop quantum corrections in the model of q -hypermultiplet interacting with vector multiplet.

We have calculated the low-energy effective action of noncommutative q -hypermultiplet in external gauge superfield. For the fundamental representation of $U(N)$ gauge group the effective action is found to be free of nonplanar contributions and UV/IR-mixing. This allows to calculate the holomorphic potential using the harmonic superspace approach and to represent it in manifestly gauge invariant form. We find that the holomorphic effective action is exactly the $\star$ -product generalization of corresponding commutative one.

For the case of adjoint representation of $U(N)$ group the situation is different. It is shown that the leading terms in the low-energy effective action are those which are responsible for the UV/IR-mixing in this model. The UV/IR-mixing appears only in $U(1)$ sector of $U(N)$ group while the $SU(N)$ part is responsible for the contributions to the holomorphic potential.

In conclusion let us briefly comment on the gauge invariance of the effective action of q -hypermultiplet in adjoint representation. At first sight the contribution (54) does not preserve the gauge symmetry since it contains the $\ln \frac{4}{m^2 p \circ p}$ factor which does not allow to restore the $\star$ -product of superfields in the effective action. It is the common problem of noncommutative gauge theories that the effective action can not be expressed in terms of $\star$ -product only but it involves the generalized $\star$ -multiplications of fields ($\star_n$ -products) [11, 15, 16]. It is clear that the effective action must be gauge invariant since we start from

the gauge invariant classical theory. In the papers [15, 16] it was shown that the gauge invariant effective action is built in terms of open Wilson line operators which absorb all the $\star_n$ -products into gauge invariant objects. Moreover, in the recent papers [27, 28] it was proved that manifestly singular logarithmic factors, like in eq. (54), do not spoil the gauge invariance but they are included into gauge invariant constructions with help of open Wilson line operators. We are faced with the same situation of logarithmic divergences in noncommutative hypermultiplet model in adjoint representation. But it is worth pointing out that this problem does not appear for the q -hypermultiplet in fundamental representation since this model is free of nonplanar diagrams.

The author is very grateful to I.L. Buchbinder for the supervision and careful reading of the paper. I would like also to thank B.M. Zupnik for valuable discussions and A. Armoni for useful comments. The work was supported in part by INTAS grant, INTAS-00-00254.

References

- [1] A. Connes, M.R. Douglas and A. Schwarz, "Noncommutative geometry and matrix theory: Compactification on tori", JHEP **9802** (1998) 003, [hep-th/9711162](#); M.R. Douglas and C. Hull, "D-branes and the noncommutative torus", JHEP **9802** (1998) 008, [hep-th/9711165](#).
- [2] N. Seiberg and E. Witten, "String theory and noncommutative geometry", JHEP **9909** (1999) 032, [hep-th/9908142](#).
- [3] M.R. Douglas, N.A. Nekrasov, "Noncommutative Field Theory", [hep-th/0106048](#); R.J. Szabo, "Quantum Field Theory on Noncommutative Space", [hep-th/0109162](#).
- [4] S. Minwalla, M. Van Raamsdonk and N. Seiberg, "Noncommutative perturbative dynamics", JHEP **0002** (2000) 020, [hep-th/9912072](#); M. Van Raamsdonk and N. Seiberg, "Comments on noncommutative perturbative dynamics", JHEP **0003** (2000) 035, [hep-th/0002186](#).
- [5] I.Ya. Aref'eva, D.M. Belov, A.S. Koshelev, Phys. Lett. **B476** (2000) 431, [hep-th/9912075](#); I.Ya. Aref'eva, D.M. Belov, A.S. Koshelev, "A note on UV/IR for noncommutative complex scalar field", [hep-th/0001215](#); I.Ya. Aref'eva, D.M. Belov, A.S. Koshelev, O.A. Rytchkov, Nucl. Phys. Proc. Suppl. **102** (2001) 11, [hep-th/0003176](#).
- [6] S. Ferrara, M.A. Lledo, "Some aspects of deformations of supersymmetric field theories", JHEP **0005** (2000) 008, [hep-th/0002084](#).
- [7] D. Klemm, S. Penati, L. Tamassia, "Non(anti)commutative superspace", [hep-th/0104190](#).
- [8] A. Armoni, Nucl. Phys. B593 (2001) 229, [hep-th/0005208](#).
- [9] A. Armoni, R. Minasian and S. Theisen, Phys. Lett. B513 (2001) 406, [hep-th/0102007](#).

- [10] T.J. Hollowood, V.V. Khoze and G. Travaglini, "Exact results in noncommutative $N = 2$ supersymmetric gauge theories", JHEP **0105** (2001) 051, [hep-th/0102045](#); C-S. Chu, V.V. Khoze and G. Travaglini, Phys. Lett. **B513** (2001) 200, [hep-th/0105187](#).
- [11] D. Zanon, Phys. Lett. **B504** (2001) 101, [hep-th/0009196](#); A. Santambrogio, D. Zanon, "One-loop four-point function in noncommutative $N = 4$ Yang-Mills theory", JHEP **0101** (2001) 024, [hep-th/0010275](#); D. Zanon, Phys. Lett. **B502** (2001) 265, [hep-th/0012009](#).
- [12] I.L. Buchbinder, I.B. Samsonov, "Noncommutative $\mathcal{N} = 2$ Supersymmetric Theories in Harmonic Superspace", [hep-th/0109130](#).
- [13] I.L. Buchbinder, M. Gomes, A.Yu. Petrov, V.O. Rivelles, Phys. Lett. **B517** (2001) 191, [hep-th/0107022](#).
- [14] A. Hasimoto and N. Itzhaki, Phys. Lett. **B456** (1999) 142, [hep-th/9907166](#).
- [15] H. Liu and J. Michelson, Nucl. Phys. **B614** (2001) 279, [hep-th/0008205](#); H. Liu, Nucl. Phys. **B614** (2001) 305, [hep-th/0011125](#).
- [16] T. Mehen, M.B. Wise, "Generalized $\ast$ -Products, Wilson Lines and the Solution of the Seiberg-Witten Equations", JHEP **0012** (2000) 008, [hep-th/0010204](#).
- [17] A. Belhaj, M. Hassaini, E.M. Sahraoui, E.H. Saidi, Class. Quant. Grav. **18** (2001) 2339, [hep-th/0007137](#).
- [18] A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky, E. Sokatchev, Class. Quant. Grav. **1**, (1984) 469; A. Galperin, E. Ivanov, V. Ogievetsky, E. Sokatchev, Class. Quant. Grav. **2** (1985) 601; 617.
- [19] A. Galperin, E. Ivanov, V. Ogievetsky, E. Sokatchev, *Harmonic Superspace*, Cambridge Univ. Press, 2001.
- [20] B. Zupnik, Sov. J. Nucl. Phys. **44** (1986) 512.
- [21] B.M. Zupnik, Phys. Lett. **B183** (1987) 175.
- [22] I.L. Buchbinder, S.M. Kuzenko, Class. Quant. Grav. **17** (1997) L157, [hep-th/9704002](#).
- [23] E.A. Ivanov, S.V. Ketov, B.M. Zupnik, Nucl. Phys. **B509** (1998) 53, [hep-th/9706078](#).
- [24] I.L. Buchbinder, E.I. Buchbinder, E.A. Ivanov, S.M. Kuzenko, B.A. Ovrut, Phys. Lett. **B412**, 309, 1997, [hep-th/9703147](#).
- [25] E.I. Buchbinder, I.L. Buchbinder, E.A. Ivanov and S.M. Kuzenko, Mod. Phys. Lett. **A13** (1998) 1071, [hep-th/9803176](#).
- [26] I.L. Buchbinder, I.B. Samsonov. Mod. Phys. Lett. **14** (1999) 2537, [hep-th/9909183](#).
- [27] M. Van Raamsdonk, "The Meaning of Infrared Singularities in Noncommutative Gauge Theories", JHEP **0111** (2001) 006, [hep-th/0110093](#).

- [28] A. Armoni, E. Lopez, "UV/IR Mixing via Closed Strings and Tachyonic Instabilities", [hep-th/0110113](#).
- [29] A.O. Barut, R. Raczka, *Theory of Group Representations and Applications*, PWN – Polish Scientific Publishers, Warszawa, 1977.